

**KRYSTEXXA (PEGLOTICASE)
INFUSION ORDERS**

****REQUIRED INFORMATION****

- This signed order form from the provider
- Patient demographics & insurance information
- Clinical/Progress Notes, Labs, Tests** supporting primary diagnosis
- Baseline Uric Acid < 6.0 mg/dl

***Patient must have Uric Acid level drawn 24-72 hours prior to infusion**

***Patient must have Glucose-6-phosphate dehydrogenase (G6PD) deficiency screening prior to initiating therapy**

Patient Name:	DOB:
Allergies:	Patient Phone:

- Diagnosis:** Chronic Gouty Arthropathy w/tophus (tophi) (ICD-10 Code: _____)
 Chronic Arthropathy w/o mention of tophus (tophi) (ICD-10 Code: _____)

J Code: J2507

KRYSTEXXA ORDERS

- Krystexxa (pegloticase) 8mg IV in 250ml of NS IV over 120 minutes

***Patient will be observed 1 hr post infusion**

Frequency: Every 2 weeks

Protocol Pre-Medication Orders: Solu-Medrol 125mg IV Antihistamine 25mg PO/IV

Patient advised to take antihistamine day before infusion

Additional Instructions:

Physician Name:	Phone:	Fax:
**Physician Signature:	Date:	